

## Second Autologous Stem Cell Implantation in Patients with Chronic Spinal Cord Injuries

Herrera AB<sup>1\*</sup> and Rodriguez PRL<sup>2</sup>

<sup>1</sup>Profesor Consultante, Profesor Titular, Especialista en Primero y Segundo Grado en Ortopedia y Traumatología, cuba

<sup>2</sup>Profesor Consultante, Profesor Auxiliar, Especialista en Primero y Segundo Grado en Cirugía General e Investigador Auxiliar, cuba

### \*Corresponding author:

Alberto Benitez Herrera,  
Profesor Consultante, Profesor Titular, Especialista  
en Primero y Segundo Grado en Ortopedia y  
Traumatología, Calle Cisneros Betancourt No. 659  
entre Bella y Aldabó. Los Pinos. Arroyo Naranjo.  
LA Habana. Cuba, Tel: +53 76443651;  
E-mail: abenith@infomed.sld.cu

Received: 15 Oct 2021

Accepted: 28 Oct 2021

Published: 02 Nov 2021

J Short Name: COS

### Copyright:

©2021 Herrera AB, This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

### Citation:

Herrera AB. Second Autologous Stem Cell Implantation in Patients with Chronic Spinal Cord Injuries. Clin Surg. 2021; 6(10): 1-2

### Keywords:

Implant; Stem cells; Regenerative medicine

### &Authors' contributions:

Herrera AB, Rodriguez PRL and all these authors are equally contributed to this work.

## 1. Abstract

**1.1. Introduction:** Since the beginning of the last century, multiple investigations have been carried out in the field of Regenerative Medicine. Stem cells obtained by different routes have been implanted in different injured tissue in order to achieve its regeneration.

**1.2. Objective:** To demonstrate the feasibility and safety of the implantation of autologous MO-CMA by subarachnoid puncture.

**1.3. Method:** A study is carried out by implanting autologous stem cells by subarachnoid puncture in five patients who had undergone a first open implant due to a neurological lesion of traumatic etiology. January 2016 to December 2018, at the Enrique Cabrera General Teaching Hospital.

**1.4. Results:** The second implant was completed in 5 patients and the following advantages were obtained: less surgical time, less possibility of sepsis, no blood loss, less hospital stay and fewer complications.

**1.5. Conclusions:** The feasibility and safety of stem cell implantation by subarachnoid puncture without complications is demonstrated in this study.

## 2. Introduction

In recent years there has been an extraordinary advance in knowledge related to different biomedical branches, including cell biology. This has given a remarkable boost to a new branch of medicine called regenerative medicine, which has been considered a "New

Revolution in Medicine". This medical discipline has been based primarily on new knowledge about stem cells and their ability to become cells of different tissues and whose objective is to stimulate or regenerate cells, tissues or organs in order to restore or establish normal function [1 -5]

## 3. Methods

We have carried out a descriptive observational cross-sectional study in thirty patients with a first open implant of autologous adult stem cells obtained from bone marrow (ACM-MO) in patients with chronic spinal cord injury of traumatic etiology with encouraging results.

We have recently practiced a second implant in 5 cases of stem cells by subarachnoid puncture, from January 2016 to 2018 at the Enrique Cabrera General Teaching Hospital.

## 4. Description of The Technique

The patient is admitted with the corresponding investigations and approval for anesthesia, two days before the implant, stimulation with leukocin (growth factor of granulocyte colonies) begins at a rate of 5 mg / kg of weights every 12 hours until 4 doses are applied.

Stem cells are obtained from peripheral blood by the centrifugation method in the laboratory.

In the operating room in a sitting position, after washing the area with soap and water, antisepsis is performed with iodinated alcohol, field cloths are placed and the intervertebral space correspond-

ing to the spinal cord injury is located, skin wheal is performed with lidocaine 1 % and then it infiltrates deeper planes.

Following the anterior scar and under radiographic control, a subarachnoid puncture with trocar # 22 is performed above and below the lesion, implanting 5cc stem cells in each intervertebral space, confirming the location of the space with the previous aspiration of cerebrospinal fluid. trocar and the rest of stem cells are administered by IV route in 100cc of physiological saline for 30 minutes.

## 5. Results

### 5.1. Patient Relationship

Patient Diagnosis 2nd Implant Complications

E.PR Spinal cord injury T-12 6/29/2016 No

MJ.GC Spinal cord injury T11-T12 10/31/2016 No

JA.OP Spinal cord injury T10-T11 11/22/2016 No

Y.GC Spinal cord injury T8-T9 3/28/2018 No

Y.MC Spinal cord injury C5-C6 11/21/2018 No

This technique has the following advantages:

Less surgical time

Less chance of sepsis.

No blood loss.

Less hospital stay

Fewer anesthetic complications.

In all cases it has been possible to carry out the procedure without great difficulties and there have been no complications.

## 6. Discussion

Regenerative Medicine is based on the potential of stem cells to become cells of different tissues [6-10].

Autologous adult stem cells used to treat spinal cord injury patients have several advantages.

First: Problems associated with immune rejection or graft-versus-host reactions can be avoided [11-13].

Second: CMA-MO transplantation is considered a safe method since its association with carcinogenesis has not been reported [14,15].

Third: There is great experience with the use of transplantation of these cells in hematological diseases [16].

## 7. Conclusions

The feasibility and safety of stem cell implantation by subarachnoid puncture is demonstrated. None of our cases have presented complications that could put this technique at risk.

## References

1. Okubo T, Iwanami A, Kohyama J. Pretreatment with a gammasecretase inhibitor prevents tumor-like overgrowth in human iPSC-derived transplants for spinal cord injury. *Stem Cell Reports*. 2016; 7: 649-63.
2. Fujimoto Y, Abematsu M, Falk A. Treatment of a mouse model of spinal cord injury by transplantation of human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem cells. *Stem Cells*. 2012; 30: 1163-73.
3. Romanyuk N, Amemori T, Turnovcova K. Beneficial effect of human induced pluripotent stem cell-derived neural precursors in spinal cord injury repair. *Cell Transplant*. 2015; 24: 1781-97.
4. Enomoto M. Therapeutic effects of neurotrophic factors in experimental spinal cord injury models. *J Neurorestoratol*. 2016; 4: 15-22.
5. Fuhrmann T, Anandakumaran PN, Shoichet MS. Combinatorial therapies after spinal cord injury: how can biomaterials help? *Adv Health Mater*. Epub. 2017; 6.
6. Raspa A, Pugliese R, Maleki M, Gelain F. Recent therapeutic approaches for spinal cord injury. *Biotechnol Bioeng*. 2016; 113: 253-9.
7. Slotkin JR, Pritchard CD, Luque B. Biodegradable scaffolds promote tissue remodeling and functional improvement in non-human primates with acute spinal cord injury. *Biomaterials*. 2017; 123: 63-76.
8. Li X, Zhao Y, Cheng S. Cetuximab modified collagen scaffold directs neurogenesis of injury-activated endogenous neural stem cells for acute spinal cord injury repair. *Biomaterials*. 2017; 137: 73-86.
9. Iyer NR, Wilems TS, Sakiyama-Elbert SE. Stem cells for spinal cord injury: strategies to inform differentiation and transplantation. *Biotechnol Bioeng*. 2017; 114: 245-59.
10. Chen C, Zhao ML, Zhang RK. Collagen / heparin sulfate scaffolds fabricated by a 3D bioprinter improved mechanical properties and neurological function after spinal cord injury in rats. *J Biomed Mater Res A*. 2017; 105: 1324-32.
11. Nejati-Koshki K, Mortazavi Y, Pilehvar-Soltanahmadi Y, Sheoran S, Zarghami N. An update on application of nanotechnology and stem cells in spinal cord injury regeneration. *Biomed Pharmacother*. 2017; 90: 85-92.
12. Nagoshi N, Okano H. Applications of induced pluripotent stem cell technologies in spinal cord injury. *J Neurochem*. Epub. 2017; 141: 848-60.
13. Karimi-Abdolrezaee S, Schut D, Wang J, Fehlings MG. Chondroitinase and growth factors enhance activation and oligodendrocyte differentiation of endogenous neural precursor cells after spinal cord injury. *PLoS One*. 2012; 7: 37589.
14. Kaneko S, Iwanami A, Nakamura M. A selective Sema3A inhibitor enhances regenerative responses and functional recovery of the injured spinal cord. *Nat Med*. 2006; 12: 1380-9.
15. US Food & Drug Administration. *Vaccines, Blood & Biologics*. 2017.
16. Hernández Ramírez Q. Is treatment with regenerative cell therapy really useful? *Rev Cubana Hematol Inmunol Hemoter* [online magazine]. 2017; 33.